



Original Research article

Ultrasonic and Microwave Effects on the Benzamide/Sulfonyl Chloride Mediated Benzoylation of Benzene Derivatives under Vilsmeier-Haack Conditions

Kamatala Chinna Rajanna*, Marri Venkateswarlu, Mukka Satish Kumar, Purugula Venkanna, Pondichery Kuppuswamy Saiprakash

Department of Chemistry, Osmania University, Hyderabad-500 007, T.S., India

ARTICLE INFO

Article history

Submitted: 2023-06-23

Revised: 2023-08-26

Accepted: 2023-09-03

Manuscript ID: CHEMM-1709-1010

Checked for Plagiarism: Yes

Language Editor:

Aghaie, Ermia

Editor who approved publication:

Dr. Zahra Abbasi

DOI: 10.48309/chemm.2023.50558

ABSTRACT

Benzamide/ Sulfonyl Chloride (SO_2Cl_2) reagent has been developed as a Vilsmeier-Haack (VH) reagent for benzoylation of aromatic compounds by using the conventional and ultrasonic sonication (US), and microwave (MW) conditions. Benzoylation is much more competent and faster than the analogous VH ((Benzamide/ POCl_3) or (Benzamide/ SOCl_2)) reagents and afforded benzoyl derivatives in good yields. Reaction times are moderately less than those observed with [Benzamide/ POCl_3] and (Benzamide/ SOCl_2) reagents. Reaction times are far less in microwave assisted (MWAR) reactions than ultrasonically assisted (USAS) reactions, which in turn are less than the conventional method with a decreasing trend: MWAR (seconds) \ll Sonication (minutes) \ll conventional (h). The highly remarkable rate accelerations in MW assisted reactions are attributed to the bulk activation of the molecules; while US assisted reactions could be accounted for cavitation effects.


KEYWORDS

Ultrasonic and microwave irradiation

Benzamide/ SO_2Cl_2

Rate accelerations

GRAPHICAL ABSTRACT



Where R = EWG or EDG

* Corresponding author: Kamatala Chinna Rajanna

E-mail: kcrajannaou@yahoo.com

© 2023 by SPC (Sami Publishing Company)

Introduction

Benzoylation forms an important class among the Friedel Crafts acylation reactions. Benzoylation reaction is generally carried out using benzoyl chloride with a Lewis acid as the benzoylating agent [1, 2]. It is a colorless fuming liquid with an irritating odor; however, it is still widely used for producing the peroxides, preparation of dyes, perfumes, pharmaceuticals, and resins. Also, it is commonly used in many laboratories because of its availability and low cost. Besides the benzoyl chloride, a number of reagents such as, benzoic anhydride, benzoyl tetrazole, 2-benzoyl-1-methylpyridinium chloride, S-benzoic-O, O-diethylphosphoro dithioic anhydride, and benzoyl cyanide, could be used for this reaction [3-6]. The reaction is usually catalyzed by the bases such as, pyridine, triethylamine, and sodium hydroxide [7-9]. Recently, Satya Paul *et al.* [8] developed a rapid, economic, and environmentally friendly method for benzoylation of -NH₂, -OH and -SH groups by using the PhCOCl-Py/basic alumina. The developed reagent system was found a good alternative to classical method since the benzoylation underwent expeditiously with high yields under the solvent-free conditions. Vasanth Chowdhury *et al.* [10] investigated the liquid phase benzoylation of benzene by benzyl chloride over the Ga₂O₃, In₂O₃, GaCl₃ and InCl₃ impregnated H_β zeolite catalysts at 60 and 80 °C as a function of reaction time has been carried out and the catalysts were compared for their acidity and catalytic performance in the benzoylation process. Benzoylation and benzoylation of different aromatic compounds (at 80 °C) over the (In₂O₃/H_β) catalyst indicated the highest benzene benzoylation activity in this study. Benzoylation is one of the most important routes for synthesis of the aromatic ketones owing to its several industrial applications ranging from petrochemicals to pharmaceuticals [1, 2]. Benzophenones are very much useful as dye intermediates, UV absorbents Also, they have immense importance in perfume industry as an additive. Recent literature reports revealed that a wide number of naturally occurring prenylated

and isoprenylated benzophenones act as potent antimicrobial, anticancer, and cytotoxic agents [3]. In addition, they are reported as new class of non-nucleoside HIV reverse transcriptase inhibitors [11-13].

On the other hand, benzoylation has also been carried out using benzamide/POCl₃, or benzamide/SOCl₂ as Vilsmeier-Haack reagents [14-17].

Over the last few years, with the increasing interest in the application of non-conventional energy sources in organic synthesis, ultrasound and microwave assisted protocols have attracted a great deal of attention from organic chemists. A great number of organic reactions have been carried out under ultrasonic and microwave irradiation to obtain higher yields, with the shorter reaction times or milder conditions [18-22]. Over the last few years, our group has been actively working on exploiting the use of eco-friendly materials such as a micelle forming surfactants and non-conventional energy sources (such as microwave and ultra sound) to serve multitudinous organic transformations [23, 24] including Vilsmeier-Haack reactions [23] and other electrophilic substitution reactions [24]. Sulfuryl chloride is a versatile and potential reagent in organic synthesis, which has been used to produce chlorinate electron-rich aryl and alkyl aryl compounds [25, 26]. Reactions with sulfuryl chloride are generally high yielding when the functionalities are compatible with this reagent. In our previous study, we have explored the use of acetamide/SO₂Cl₂ as an effective reagent for acylation of aromatic compounds under ultrasonic and microwave conditions [27]. Encouraged by the results presented therein, we have embarked on the present study to explore the use of benzamide/SO₂Cl₂ as a new VH reagent for effective benzoylation of aromatic compounds under conventional and nonconventional conditions. The present developed reagent system is found to be a good alternative to classical method since the benzoylation underwent expeditiously with high yields under the solvent-free conditions.

Experimental

All the chemicals were purchased from Aldrich or Arcos Organics and used without further purification. Analytical TLC was carried out using Merck aluminum- backed 0.2 mm silica gel 60 F-254 plates. Column chromatography was conducted using Merck silica gel 60 (230-400mesh). Ultrasonically assisted reactions were performed in a Sonicator bath (KQ-250B, China). A flat transducer with a frequency of 40 kHz and voltage of 220 V (with an output of 100 W electric power rating) was mounted at the bottom of the Sonicator.

Preparation of benzamide/SO₂Cl₂ (Sulfuryl Chloride) reagent

Requisite amounts (equimolar) of benzamide and sulfuryl chloride (SO₂Cl₂) are mixed in acetonitrile under chilled (at -5°C) conditions for the preparation of benzamide/SO₂Cl₂ (sulfuryl chloride) reagent. The reagent is always prepared afresh before use.

Representative procedure for benzylation using SO₂Cl₂/benzamide reagent under reflux condition

About 0.01 mol of aromatic substrate and 0.015 mol of SO₂Cl₂/benzamide reagent and acetonitrile solvent were taken in a clean Round bottom flask and refluxed for about 5 to 8 hours and progress of the reaction was monitored by checked by thin layer chromatography. After completion, the reaction mixture is washed with NaHCO₃ solution, followed by the addition of ethyl acetate. The organic layer was separated, dried over Na₂SO₄, under the vacuum to get pure product. The purified products were characterized by ¹HNMR and mass spectroscopic technique and also by their physical data, which were similar to our earlier reports [14].

Representative procedure for benzylation using of SO₂Cl₂/benzamide reagent under sonication

Centimolar (0.01 mol) organic substrate, SO₂Cl₂/benzamide reagent (0.015 mol) and acetonitrile were added in a previously cleaned in a round bottom flask and clamped in a sonicator and progress of the reaction is followed by thin

layer chromatography. After completion of the reaction, product is separated as described in the preceding section.


Representative procedure for microwave assisted benzylation using of SO₂Cl₂/benzamide reagent under microwave irradiation

A centimolar (0.01 mol) aromatic compound, about 0.015 mol of SO₂Cl₂/benzamide, were taken in a cleaned in a Round bottom flask containing acetonitrile and clamped in a laboratory MW oven. Progress of the reaction is monitored by thin layer chromatography. The separation and purification process is by and large similar to the above procedure.

Results and Discussion

Freshly prepared benzamide/SO₂Cl₂ was used for the benzylation of aromatic compounds under conventional, ultrasonic, and microwave assisted conditions.

Aromatic compounds underwent the smooth benzylation and afforded very good yields of products when treated with benzamide/SO₂Cl₂, respectively under the conventional and non-conventional conditions (Scheme 1). The reaction times observed with benzamide/SO₂Cl₂ (Sulfuryl chloride) adduct are in the range of 5-8 h These reaction times observed in this study are much less than those observed with benzamide/(thionyl chloride) SOCl₂ (10-13h) or benzamide/POCl₃ (11-13 h) when used as benzylation agents in our previous study [13]. This trend can be probably attributed to the increased Lewis acid character of SO₂Cl₂ (Sulfuryl chloride) over (thionyl chloride) SOCl₂ or POCl₃ which favors the formation of a better iminium salt with benzamide than that of the latter oxychlorides. We have successfully extended our study conducted under ultrasonically and microwave assisted conditions. The results are summarized in Table 1. Mechanism of the reaction could be explained through the formation of iminium salt type intermediate in the first step due to interaction of benzamide and sulfuryl chloride, which then converts aromatic compound to benzoyl derivatives followed by the removal of HCl and H₂SO₄ in the fast steps as



Scheme 3: Mechanism of benzoylation of aromatic compounds using benzamide/ SO_2Cl_2

Reaction times reduced from 5-8 h (under conventional methods) to 40-60 min (under sonication) and only 4-6 min under microwave assisted conditions. It is earlier reported that sonication of a chemical reaction in solution triggers the formation, growth, and collapse of bubbles, called as acoustic cavitation [17-22]. During sonication, compression of the liquid is followed by the rarefaction (expansion), in which a sudden pressure drop forms small oscillating bubbles, which expand with each cycle of the applied ultrasonic energy until they reach an unstable size; they can then collide and/or violently collapse. This bubble collapse causes intense local heating, and high-pressures which are responsible for rate enhancements and shorter reaction times. Acoustic cavitation effects could thus explain the experimental results in this study. On the other hand, microwave irradiation causes enhanced dipole-dipole and ionic interactions apart from direct absorption of

the selective microwave energy by the reactive species irrespective of their polar or dipolar nature. This may lead to the bulk activation of molecules, which decompose or rearrange to afford the products and bring about highly significant decrease in reaction times and increase in the product yield. Reactions with large activation energies will also be influenced greatly with microwave irradiation. According to Haile *et al.* [28], microwave assisted reactions get advantages such as volumetric heating, selectivity, rapid kinetics, homogeneity, and energy saving due to the direct absorption of the selective microwave energy. Microwave irradiation also provides pollution free environment since there are no byproducts of combustion. Thus, extremely faster reaction rates in MIR system could be accounted for the bulk activation of molecules rather than random activation.

Table 1: benzylation of aromatic compounds under different conditions

Entry	Substrate	Product	Conventional		Sonication		Microwave	
			Time (h)	Yield (%)	Time (min)	Yield (%)	Time (min)	Yield (%)
1	Toulene	4-Me benzophenone	5	72	45	74	4	70
2	Chlorobenzene	4-Cl benzophenone	4.5	70	40	72	4	70
3	Nitrobenzene	4-NO ₂ benzophenone	4	75	35	78	3.5	73
4	Fluorobenzene	4-F benzophenone	5	71	50	74	5	72
5	Anisole	4-OMe benzophenone	6	74	60	76	5	75
6	Phenol	4-OH benzophenone	6.5	75	70	77	4.5	78
7	Bromo Benzene	4-Br benzophenone	4	68	40	70	4	74
8	Naphthalene	2-Benzoyl naphthalene	7.5	65	65	69	6	66
9	Acetophenone	3-Benzoyl acetophenone	7	76	60	80	5	80
10	4-OH acetophenone	3-Benzoyl 4-OH acetophenone	4	76	40	78	4.5	82
11	4-Cl acetophenone	3-Benzoyl 4-Cl acetophenone	5.5	62	45	67	4	69
12	4-NO ₂ acetophenone	3-Benzoyl 4-NO ₂ acetophenone	4	79	50	80	3.5	82
13	Benzaldehyde	3-Benzoyl benzaldehyde	7.5	70	60	75	5	78
14	4-Cl benzaldehyde	3-Benzoyl 4-Cl benzaldehyde	5.5	74	40	76	4	81
15	4-OMe benzaldehyde	3-Benzoyl 4-OMe benzaldehyde	6.5	70	50	75	5.5	79

Data presented in Table 1 shows that benzylation occurred exclusively at the para position to the substituents such as -OMe, -Me, -Cl and -Br in mono substituted benzenes and afforded quantitative yields. However, in cases where the para positions are blocked (Table 1, entries 10,11,12,14 and 15); the benzoyl group is introduced in the *meta* position (3-position). The benzylation of naphthalene is observed at second position (2 position) of naphthalene with good yield (Table 1, entry 8). On the whole benzylation took place efficiently with good regioselectivity in normal as well as ultrasound and microwave assisted reactions.

Conclusions

In summary, we have successfully developed a new protocol for benzylation of aromatic compounds under conventional, sonication and microwave irradiation. Longer reaction times (5-8 hours) under the normal conditions reduced to 40-60 min under sonication, while microwave assisted reactions further reduced the reaction time to only 4-6 minutes. Reactions are conducted with economically cheap and readily available laboratory desktop chemicals with a

simple work. Benzylation occurred with good regioselectivity. Rate enhancements coupled with enhanced reaction yields under sonication and microwave irradiation substantiate that the present work is a good contribution in the area of benzylation reactions of aromatic compounds with potential biological activity.

Acknowledgment

The authors would like to thank UGC and New Delhi for awarding Rajiv Gandhi Post Doc fellowship to Dr. M.V. Authours sincerely acknowledge their gratitude to Prof. T. Navaneeth Rao (Former Vice-Chancellor, O.U) for constant encouragement.

Disclosure Statement

No potential conflict of interest was reported by the authors.

Funding

This study did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

All authors contributed toward data analysis, drafting, and revising the paper and agreed to responsible for all the aspects of this work.

Conflict of interest

The authors declare that they have no conflicts of interest in this article.

References

- [1]. Pearson A.g., Roush W.J., *Activating Agents and Protecting Groups*, 1999 [[Google Scholar](#)], [[Publisher](#)]
- [2]. (a) Olah G.A., Friedel–Crafts and Related Reactions, 1964 [[Google Scholar](#)], [[Publisher](#)] (b) Olah G.A., *Friedel– Crafts Chemistry*, Wiley-Interscience, New York, USA, 1973, III–IV [[Publisher](#)]
- [3]. Clarke H.T., Rahrs E.J., *Organic Syntheses. Collective Volumes, Coll*, 1941, 1:91 [[Google Scholar](#)]
- [4]. Stawinski J., Hozumi T., Narang S.A., Benzoyltetrazole: a mild benzoylating reagent for nucleosides, *Journal of the Chemical Society, Chemical Communications*, 1976, 3:243 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [5]. Yamada M., Watabe Y., Sakakibara T., Sudoh R., Yamada, M., Watabe, Y., Sakakibara, T. and Sudoh, R., Preparation of a water-soluble acylating agent: benzoylation of acids, amines, and phenols with 2-benzoylthio-1-methylpyridinium chloride in aqueous phase, *Journal of the Chemical Society, Chemical Communications*, 1979, 179 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6]. Carey F.A., Hodgson K.O., Efficient syntheses of methyl 2-O-benzoyl-4, 6-O-benzylidene- α -D-glucopyranoside and methyl 2-O-benzoyl-4, 6-O-benzylidene- α -D-ribo-hexopyranosid-3-ulose, *Carbohydrate Research*, 1970, 12:463 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7]. Greene T.W., Wuts P.G., *Protective groups in organic synthesis*, 1999 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [8]. McOmie J.F.W. ed., *Protective Groups in Organic Chemistry*, 1973 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [9]. Paul S., Nanda P., Gupta R., PhCOCl-Py/basic alumina as a versatile reagent for benzoylation in solvent-free conditions, *Molecules*, 2003, 8b374 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [10]. Choudhary V.R., Jana S.K., Patil N.S., Bhargava S.K., Friedel–Crafts type benzoylation and benzoylation of aromatic compounds over H β zeolite modified by oxides or chlorides of gallium and indium, *Microporous and mesoporous materials*, 2003, 57:21 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [11]. Pecchio M., Solis P.N., Lopez-Perez J.L., Vsquez Y., Rodriguez N., Olmedo D., Correa M., Feliciano A., Gupta M.P., Cytotoxic and antimicrobial benzophenones from the leaves of *Tovomita longifolia*. *Journal of natural products*, 2006, 69:410 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [12]. Ren J., Chamberlain P.P., Stamp A., Short S.A., Weaver K.L., Romines K.R., Hazen R., Freeman A., Ferris R.G., Andrews C.W., Boone L., Structural basis for the improved drug resistance profile of new generation benzophenone non-nucleoside HIV-1 reverse transcriptase inhibitors, *Journal of medicinal chemistry*, 2008, 51:5000 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [13]. Geetha P.K., Setty M.B., Baswaraj B., Comparison Of The Solubility Of Luting Cements Immersed In Artificial Saliva-An Invitro Study. *Indian Journal of dental sciences*, 2011, 3 [[Google Scholar](#)]
- [14]. Rajanna K.C., Venkateswarlu M., Satish Kumar M., Umesh Kumar U., Venkateswarlu G., Saiprakash P.K., Kinetics and mechanism of certain benzoylation reactions under Vilsmeier–Haack conditions using benzamide and oxychloride in acetonitrile medium, *International Journal of Chemical Kinetics*, 2013, 45:69 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [15]. (a) Martin G.J., Poignant S. , Filleux M.L., Quemeneuer M.T., Recherches sur la reaction de vilsmeier-haack etude du mecanisme de formation du complexe par des mesures cinetiques en resonance magnetique nucleaire. *Tetrahedron Letters*, 1970, 11:5061 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [16]. Katritzky A.R., Meth-Cohn O., Rees C.W. eds., 1995. *Comprehensive Organic Functional Group*

- Transformations: Synthesis: carbon with one heteroatom attached by a single bond.* Elsevier. [[Google Scholar](#)], [[Publisher](#)]
- [17]. Davis T.L., Yelland W.E., Addition of butylamine to butyl isocyanide, *Journal of the American Chemical Society*, 1937, **59**:1998 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [18]. (a) Suslick K.S. In *Kirk-Othmer Encyclopedia of Chemical Technology*; 4th Ed. J. Wiley & Sons: New York, 1998, **26**:517 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)] (b) Mason T.J., Lorimer J.P., Sonochemistry, Part 1—the physical aspects, *Chemical Society Reviews*, 1987, **16**:239 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)] (c) Einhorn C., Einhorn J., Luche J.L., Sonochemistry—The use of ultrasonic waves in synthetic organic chemistry, *Synthesis*, 1989, **1989**:787 [[Google Scholar](#)], [[Publisher](#)] (d) van Eldik R., Hubbard C.D. eds., Chemistry under extreme and non-classical conditions, *John Wiley & Sons*, 1997 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [19]. Memarian H.R., Mohammadpoor-Baltork I., Nikoofar K., Ultrasound-assisted thiocyanation of aromatic and heteroaromatic compounds using ammonium thiocyanate and DDQ, *Ultrasonics sonochemistry*, 2008, **15**:456 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20]. (a) Caddick S., Microwave assisted organic reactions, *Tetrahedron*, 1995, **51**:10403; [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)] (b) Lidström P., Tierney J., Wathey B., Westman J., Microwave assisted organic synthesis—a review, *Tetrahedron*, 2001, **57**:9225 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21]. Hayes B.L., Recent advances in microwave-assisted synthesis. *Aldrichimica Acta*, 2004, **37**:66 [[Google Scholar](#)], [[Publisher](#)] (b) Roberts B.A., Strauss C.R., Toward rapid, “green”, predictable microwave-assisted synthesis, *Accounts of Chemical Research*, 2005, **38**:653 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [22]. De la Hoz A., Diaz-Ortiz A., Moreno A., Microwaves in organic synthesis. Thermal and non-thermal microwave effects, *Chemical Society Reviews*, 200, **34**:164 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23]. (a) Ali M.M., Tasneem K.C., Rajanna P.K., . An efficient and facile synthesis of 2-chloro-3-formyl quinolines from acetanilides in micellar media by Vilsmeier-Haack cyclisation, *Synlett*, 2001, **2001**:251 [[Google Scholar](#)], [[Publisher](#)] (b) Rajanna K.C., Moazzam Ali, Sana M., Tasneem S., Saiprakash P.K., Vilsmeier Haack Acetylation in Micellar Media: An Efficient One Pot Synthesis of 2-Chloro-3-acetyl Quinolines, *Journal of dispersion science and technology*, 2004, **25**:17 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [24]. (a) Elming N., Clauson-Kaas N., Anderson E.P., Eliasson N.A., Thorell B., Transformation of 2-(Hydroxymethyl)-5-(aminomethyl)-furan into 6-Methyl-3-pyridinol, *Acta Chemica Scandinavica*, 1956, **10**:1603 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)] (b) Mavoungou-Gomes L., Cabares, J., Preparation of 5-alkyl-and 5-arylalkyl-2-methylbenzofuran-6, 7-dicarboxylic acids. Conversion of 5-arylalkylbenzofuran anhydrides to new heterocycles. *Bulletin de la Société Chimique de France*, 1986. **3**:401 [[Google Scholar](#)]
- [25]. (a) Watson W.D., Regioselective para-chlorination of activated aromatic compounds. *The Journal of Organic Chemistry*, 1985, **50**:2145 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)] (b) Smith K., Tzimas M., Brown C.M., Payne K., Dialkyl sulfides as selective catalysts for the chlorination of phenols. *Sulfur Letters*, 1999, **22**:89 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)] (c) Reiter L., Berg G.E., The synthesis of spirocyclic [1, 2, 4] triazolo [4, 3-a] quinolines as potential ligands for the benzodiazepine receptor. *Heterocycles (Sendai)*, 1992, **34**:771 [[Google Scholar](#)], [[Publisher](#)] (d) Brown H.C., Sulfuryl chloride in organic chemistry, *Industrial & Engineering Chemistry*, 1944, **36**:785 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26]. Yu G., Mason H.J., Wu X., Endo M., Douglas J., Macor J.E., A mild and efficient method for aromatic chlorination of electron-rich arylalkyl amines, *Tetrahedron Letters*, 2001, **42**:3247 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [27]. Satish Kumar M., Rajanna K.C., Venkanna P., Venkateswarlu M., Acetamide/SO₂Cl₂ as an efficient reagent for Friedel–Craft’s acylation of aromatic compounds under ultrasonic and microwave conditions, *Tetrahedron Letters*, 2014, **55**:1756 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[28]. Haile S.M., Johnson D.W., Wiseman G.H., Bowen H.K., Aqueous precipitation of spherical zinc oxide powders for varistor applications, *Journal of the American Ceramic Society*, 2004, **72**:1989 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

HOW TO CITE THIS ARTICLE

Kamatala Chinna Rajanna*, Marri Venkateswarlu, Mukka Satish Kumar, Purugula Venkanna, Pondichery Kuppaswamy Saiprakash. Ultrasonic and Microwave Effects on the Benzamide/Sulfuryl Chloride Mediated Benzoylation of Benzene Derivatives under Vilsmeier-Haack Conditions. *Chem. Methodol.*, 2023, 7(9) 650-658

DOI: <https://doi.org/10.48309/chemm.2023.50558>

URL: https://www.chemmethod.com/article_50558.html